



UNIVERSITY OF ILLINOIS
EXTENSION

**GROWING A NEW GENERATION
OF ILLINOIS FRUIT AND VEGETABLE FARMERS**

USDA NIFA Beginning Farmer and Rancher Development Program
Grant # 2012-49400-19565

INTRODUCTION TO SOILS

J. D. Kindhart & Ellen Philips
March 2015




Soil

- This definition is from Soil Taxonomy, second edition
- Soil is a natural body comprised of solids (minerals and organic matter), liquid, and gases that occurs on the land surface, occupies space, and is characterized by one or both of the following: horizons, or layers, that are distinguishable from the initial material as a result of additions, losses, transfers, and transformations of energy and matter or the ability to support rooted plants in a natural environment.

Ideal Soil Profile

0 feet - surface




2 inches to 5+ feet


Soil Horizons

- (O horizon)
- **A horizon**
 - Topsoil
- (E horizon)
- **B horizon**
 - Subsoil
- **C horizon**
 - Substratum
 - Parent Material
 - Regolith
 - Bedrock






Soil Texture



UNIVERSITY OF ILLINOIS
EXTENSION



Cation Exchange Capacity: CEC



UNIVERSITY OF ILLINOIS
EXTENSION



16 Essential Nutrients

- **Structural Nutrients**

- **Carbon (C)**
- **Oxygen (O)**
- **Hydrogen (H)**

- **Major (macro) nutrients**

- Nitrogen (N)
- Phosphorus (P)
- Potassium (K)


- **Secondary nutrients**


- **Calcium (Ca)**
- **Magnesium (Mg)**
- **Sulfur (S)**

- **Micronutrients**

- Boron (B)
- Chloride (Cl)
- Copper (Cu)
- Iron (Fe)
- Manganese (Mn)
- Molybdenum (Mo)
- Zinc (Zn)







Cation Exchange Capacity (CEC)

- The total number of exchangeable cations a soil can hold
 - amount of its negative charge

Soil Texture

CEC Range

(meq/100 g soil)

organic soils

> 50

clays

25-50

silts

8-30

sands

5-15



Figure 8.7. Cation-exchange capacity of Illinois soils. The darkest areas are sands with low capacity.

What Soil is on Your Property?

Web Soil Survey - Home - Windows Internet Explorer

http://websoilsurvey.nrcs.usda.gov/app/HomePage.htm

File Edit View Favorites Tools Help

WEB SEARCH

Web Soil Survey - Home

USDA United States Department of Agriculture Natural Resources Conservation Service

Web Soil Survey

Home About Soils Help Contact Us

You are here: Web Soil Survey Home

Search

Enter Keywords

All NRCS Sites

Browse by Subject

- Soils Home
- National Cooperative Soil Survey (NCSS)
- Archived Soil Surveys
- Status Maps
- Official Soil Series Descriptions (OSD)
- Soil Series Extent Mapping Tool
- Soil Data Mart
- Geospatial Data Gateway
- eFOTG
- National Soil Characterization Data
- Soil Geochemistry Spatial Database
- Soil Quality
- Soil Geography
- Geospatial One Stop


The simple yet powerful way to access and use soil data.

START WSS

Welcome to Web Soil Survey (WSS)

Web Soil Survey (WSS) provides soil data and information produced by the National Cooperative Soil Survey. It is operated by the USDA Natural Resources Conservation Service (NRCS) and provides access to the largest natural resource information system in the world. NRCS has soil maps and data available online for more than 95 percent of the nation's counties and anticipates having 100 percent in the near future. The site is updated and maintained online as the single authoritative source of soil survey information.

Three Basic Steps

- 1 Define...**
Area of Interest (AOI) Use the Area of Interest tab to define your area of interest.

Click to view larger image.
- 2 View/Explore...**
Soil Map Click the Soil Map tab to view or print a soil map, or click the Soil Data Explorer tab to access soil data for your

I Want To...

- Start Web Soil Survey (WSS)
- Know the requirements for running Web Soil Survey
- Know whether Web Soil Survey works in my web browser
- Know the Web Soil Survey hours of operation
- Find what areas of the U.S. have soil data

Announcements/Events

- Web Soil Survey Release History

I Want Help With...

- How to use Web Soil Survey
- How to use Web Soil Survey Online Help
- Known Problems and Workarounds
- Frequently Asked Questions
- Citing Web Soil Survey as a source of soils data

Tips & Shortcuts WSS

<http://websoilsurvey.nrcs.usda.gov/app/>



http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx

File Edit View Favorites Tools Help

Google Search Share More >> Sign In

Dupage County Deals Website Prices and Quote... Six Figure Moms At Home LifeMart Colby Digs Soil For those ...

USDA United States Department of Agriculture Natural Resources Conservation Service

Web Soil Survey

Contact Us Download Soils Data Archived Soil Surveys Soil Survey Status Glossary Preferences Link Logout Help

Area of Interest (AOI) Soil Map Soil Data Explorer Shopping Cart (Free)

Search

Area of Interest

Import AOI

Quick Navigation

Address

View ?

Address 1350 West Prairie Drive, sycamore, illinois

Show location marker ☒

View

State and County

Soil Survey Area

Latitude and Longitude

PLSS (Section, Township, Range)

Bureau of Land Management

Department of Defense

Forest Service

National Park Service

Hydrologic Unit

Area of Interest Interactive Map

Legend

View Extent Contiguous U.S. Scale (not to scale)

Mercantile Dr

Mediteranean Dr

Peace Rd

Praine Dr

Alton Rd

Bethany Rd

DeKalb

IL

0 867ft

http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx

File Edit View Favorites Tools Help

Google Search Share More >> Sign In

Dupage County Deals Website Prices and Quote... Six Figure Moms At Home LifeMart Colby Digs Soil For those ...

USDA United States Department of Agriculture Natural Resources Conservation Service

Web Soil Survey

Contact Us Download Soils Data Archived Soil Surveys Soil Survey Status Glossary Preferences Link Logout Help

Area of Interest (AOI) Soil Map Soil Data Explorer Shopping Cart (Free)

Search

Area of Interest

Open All Close All

AOI Properties

Clear AOI

AOI Information

Name

Map Unit Symbols

☒ Use Soil Survey Area Map Unit Symbols

☐ Use National Map Unit Symbols

Area (acres) 14.0

Soil Data Available from Web Soil Survey

DeKalb County, Illinois (IL037)

Spatial Data Version 2, Feb 12, 2010

Tabular Data Version 6, Jan 20, 2012

Clear AOI

Import AOI

Export AOI

Area of Interest Interactive Map

Legend

View Extent Contiguous U.S. Scale (not to scale)

DeKalb IL

Peace Rd

Bethany Rd

Prairie Dr


0 357ft

Search

Map Unit Legend

DeKalb County, Illinois (IL037)

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
154A	Flanagan silt loam, 0 to 2 percent slopes	6.9	49.0%
512B	Danabrook silt loam, 2 to 5 percent slopes	7.1	51.0%
Totals for Area of Interest		14.0	100.0%



Warning: Soil Map may not be valid at this scale.

You have zoomed in beyond the scale at which the soil map for this area is intended to be used. Mapping of soils is done at a particular scale. The soil surveys that comprise your AOI were mapped at 1:12,000. The design of map units and the level of detail shown in the resulting soil map are dependent on

DeKalb County, Illinois

Map symbol and soil name	Depth	Cation-exchange capacity	Effective cation-exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	<i>In</i>	<i>meq/100g</i>	<i>meq/100g</i>	<i>pH</i>	<i>Pct</i>	<i>Pct</i>	<i>dS/m</i>	
154A—Flanagan silt loam, 0 to 2 percent slopes								
Flanagan	0-18	17-23	—	5.6-7.3	0	0	0	0
	18-38	26-31	—	5.6-7.3	0	0	0	0
	38-45	19-26	—	5.6-7.3	0	0	0	0
	45-49	15-21	—	6.1-7.8	0-10	0	0	0
	49-60	8.5-21	—	7.4-8.4	15-40	0	0	0
512B—Danabrook silt loam, 2 to 5 percent slopes								
Danabrook	0-13	16-23	—	5.6-7.3	0	0	0	0
	13-33	19-28	—	5.1-7.3	0	0	0	0
	33-50	11-18	—	5.6-7.8	0-20	0	0	0
	50-60	7.9-11	—	7.4-8.4	15-40	0	0	0

Description — Chemical Soil Properties

Chemical Soil Properties

This table shows estimates of some chemical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable cations that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. The ability to retain cations reduces the hazard of ground-water pollution.

Effective cation-exchange capacity refers to the sum of extractable cations plus aluminum expressed in terms of milliequivalents per 100 grams of soil. It is determined for soils that have pH of less than 5.5.

Some Factors Influenced by Soil Type

- Plant Selection: grow adapted varieties
- Irrigation: Soil types directly impacts plant available water
- Drainage
- Fertilizer recommendation
- Yield potential
- Herbicides: read label for special instructions

SOIL STRUCTURE




Relationship between structure and porosity



Granular structure




Blocky structure



High residue and cover crops contribute organic matter to soil, while no-till management helps protect organic matter and allow accumulation. Organic matter provides food for earthworms and other soil biota. All play a role in developing or protecting soil structure and macropores to help soil function at a high level. Inset shows relationship of macro- and micropores to soil aggregates.

Bulk Density



Particle Density

100% solid
Weight = 2.66 g
Volume = 1 cm³

Bulk Density


50% solid, 50% pore space
Weight = 1.33 g
Volume = 1 cm³



UNIVERSITY OF ILLINOIS
EXTENSION




Penetrometer



Compacted Zone

Impermeable Layer



UNIVERSITY OF ILLINOIS
EXTENSION




Preventing and Repairing Compaction

- Break up compacted layer
 - Double digging
 - Sub soiling
 - Core aeration of lawns
- Plant deep rooted plants
 - Tillage radish
 - Annual Ryegrass

Preventing and Repairing Compaction

- Prevent further compaction
 - Minimize Tillage
 - Minimize Rototilling
 - Vary depth of tillage each year
 - Minimize walking on soil
 - Raised beds guide traffic
 - Do not drive cars or other vehicles across soil
 - Develop drive aisles



Interpreting Soil Test Results

REPORT NUMBER

REPORT DATE

RECEIVED DATE

ACCOUNT



13611 "B" Street • Omaha, Nebraska 68144-3693 • (402) 334-7770 • FAX (402) 334-9121
www.midwestlabs.com
IDENTIFICATION

PAGE 1/1

AGPRO SYSTEMS INC
DON MISCHNICK/D KINSEY
242 PR 3341
BIG SANDY TX 75755-4658

SOIL ANALYSIS REPORT

LAB NUMBER	SAMPLE IDENTIFICATION	ORGANIC MATTER MODIFIED LOI percent RATE	PHOSPHORUS			POTASSIUM		MAGNESIUM		CALCIUM		SODIUM		pH		CATION EXCHANGE CAPACITY C.E.C. meq/100g	PERCENT BASE SATURATION (COMPUTED)				
			P ₁ OMEX 1:7 ppm RATE	P ₂ STRONG BRAY 1:7 ppm RATE	BICARBONATE P ppm RATE	K ppm RATE	Mg ppm RATE	Ca ppm RATE	Na ppm RATE	SOL pH 5:1	BUFFER INDEX	% K	% Mg	% Ca	% H		% Na				
205		1		2		3		4		5		6	7	8			9				

LAB NUMBER	NITRATE-N (FIA)									SULFUR S %CAP ppm RATE	ZINC Zn DTPA ppm RATE	MANGANESE Mn DTPA ppm RATE	IRON Fe DTPA ppm RATE	COPPER Cu DTPA ppm RATE	BORON B BORL DTPA ppm RATE	SOLUBLE SALTS 1:1 method/ cm RATE	
	SURFACE			SUBSOIL 1			SUBSOIL 2										Total lb/A
205	ppm	lb/A	depth (in)	ppm	lb/A	depth (in)	ppm	lb/A	depth (in)								
										12	13	14	15	16	17	18	19

REV. 12/03

The above analytical results apply only to the sample(s) submitted. Samples are retained a maximum of 30 days.

Our reports and letters are for the exclusive and confidential use of our clients and may not be reproduced in whole or in part, nor may any reference be made



FIELD INFORMATION			
Field ID	Hill top field	Sample no.	1
Acres	40	Last Lined	Not known
		Irrigated	No
Last crop	019 Cool-Season Grass Pasture		

This report is for:

 Example Report
University of Missouri
Columbia, MO 65211

A

Serial no. M9999	Lab no. 9969999
Area 015	County 010 Region 3
Submitted	Processed
06/10/96	06/12/96

Soil sample submitted by:

B SOIL TEST INFORMATION			C RATING					
			Very low	Low	Medium	High	Very High	Excess
pH _s	(salt pH)	4.9	*****					
Phosphorus	(P)	22 lbs/acre	*****					
Potassium	(K)	303 lbs/acre	*****					
Calcium	(Ca)	2091 lbs/acre	*****					
Magnesium	(Mg)	278 lbs/acre	*****					
Sulfur	(SO ₄ -S)	ppm						
Zinc	(Zn)	ppm						
Manganese	(Mn)	ppm						
Iron	(Fe)	ppm						
Copper	(Cu)	ppm						
Organic matter	2.2 %	Neutralizable acidity	6.0	meq/100g	Cation Exch. Capacity	12.8	meq/100g	
pH in water		Electrical Conductivity		mmho/cm	Sodium (Na)		lbs/s	
Nitrate (NO ₃ -N)	Topsoil	ppm	Subsoil	ppm	Sampling Depth	Top	Inches	Subsoil
							Inches	
D NUTRIENT REQUIREMENTS								
E Cropping options	F Yield goal	Pounds per acre					G	H LIMESTONE SUGGESTIONS
		N	P ₂ O ₅	K ₂ O	Zn	S		
Alfalfa/Grass Establishment	0	20	55	0				Effective neutralizing material (ENM)
Clover/Grass Establishment	0	20	45	0				
Alfalfa/Grass Hay	6	0	80	235				Effective magnesium (EMg)
Cool-Season Grass Pasture	150 CD/A	90	30	20				

I

To determine limestone needs in tons/acre, divide ENM requirements by the guarantee of your limestone dealer.

When N requirement for cool-season grass exceeds 90 lbs/acre, apply 2/3 of it during the period from December through February, and the remainder in August.

Do not use nitrogen on spring seedlings of legumes after May 1st because of potential weed competition.

 Area Agronomy Specialist Agronomy Specialist Phone (573) 882-1000
White - Farmer, Yellow - ASOS, Blue - Firm, Pink - Extension

NP 189 Revised 1/95

Signature


University of Missouri, Lincoln University, U.S. Department of Agriculture & Local University Extension Councils Cooperating

equal opportunity institutions

$$\text{Pounds/A} = \text{ppm} \times 2$$

Acre Furrow Slice

Tilled Soil




Recommendations




Soil pH

- Measure of alkalinity or acidity on a scale of 0 - 14



Adjusting pH






Table 8.3. Suggested limestone rates based on soil type, pH, cropping system, and 9-inch depth of tillage.

Soil type ^a	Soil pH value																					
	4.5	4.6	4.7	4.8	4.9	5.0	5.1	5.2	5.3	5.4	5.5	5.6	5.7	5.8	5.9	6.0	6.1	6.2	6.3	6.4	6.5	7.0
Tons of typical limestone ^b to apply to grain farming systems																						
A	8.0	8.0	8.0	8.0	8.0	8.0	7.8	7.0	6.3	5.5	4.8	4.0	3.3	2.5	1.8	1.0	Optional					
B	8.0	8.0	7.5	7.0	6.5	6.0	5.5	5.0	4.5	4.0	3.5	3.0	2.5	2.0	1.5	1.0	Optional					
C	6.6	6.3	5.9	5.5	5.1	4.8	4.4	4.0	3.6	3.3	2.9	2.5	2.1	1.8	1.4	1.0	Optional					
D	4.0	3.8	3.6	3.4	3.2	3.0	2.8	2.6	2.4	2.2	2.0	1.8	1.6	1.4	1.2	1.0	Optional					
E	4.0	3.6	3.2	2.8	2.4	2.0																
Tons of typical limestone ^b to apply to forage farming systems (alfalfa, clover, lespedeza)																						
A	11.0	11.0	11.0	11.0	11.0	11.0	11.0	10.3	9.6	8.9	8.1	7.4	6.7	6.0	5.3	4.6	3.9	3.1	2.4	1.7	1.0	Optional
B	11.0	11.0	11.0	10.4	9.9	9.3	8.8	8.2	7.7	7.1	6.6	6.0	5.4	4.9	4.3	3.8	3.2	2.7	2.1	1.6	1.0	Optional
C	10.0	9.6	9.1	8.7	8.2	7.8	7.3	6.9	6.4	6.0	5.5	5.1	4.6	4.2	3.7	3.3	2.8	2.4	1.9	1.5	1.0	Optional
D	6.0	5.8	5.5	5.3	5.0	4.8	4.5	4.3	4.0	3.8	3.5	3.3	3.0	2.8	2.5	2.3	2.0	1.8	1.5	1.3	1.0	Optional
E	6.0	5.4	4.9	4.3	3.8	3.2	2.7	2.1	1.6	1.0												

Note: If plowing is less than 9 in., reduce the amount; if it is more than 9 in., increase it. A chisel plow, disk, or field cultivator rather than a moldboard plow may not mix limestone deeper than 4 to 5 in.; for no-till or pasture systems, use the equivalent of a 3-in. tillage depth (one-third of the amount suggested).

^aSoil A: Dark-colored silty clays and silty clay loams (CEC > 24). Soil B: Light- and medium-colored silty clays and silty clay loams; dark-colored silt and clay loams (CEC 15–24). Soil C: Light- and medium-colored silt and clay loams; dark- and medium-colored loams; dark-colored sandy loams (CEC 8–15). Soil D: Light-colored loams; light- and medium-colored sandy loams; sands (CEC < 8). Soil E: Muck and peat. Soil color is usually related to organic matter. Light-colored soils <2.5% organic matter; medium-colored soils 2.5–4.5% organic matter; dark-colored soils >4.5% organic matter.

^bTypical limestone: 10% of the particles are greater than 8-mesh; 30% pass an 8-mesh and are held on 30-mesh; 30% pass a 30-mesh and are held on 60-mesh; and 30% pass a 60-mesh. A calcium carbonate equivalent (total neutralizing power) of 90%. Effective neutralizing value (ENV) of this material is 46.35 for 1 year after application, and 67.5 for 4 years after application. To correct the rate of application based on the ENV of the material available, follow calculations in the worksheet on page 98.



COOPERATIVE EXTENSION SERVICE
UNIVERSITY OF KENTUCKY COLLEGE OF AGRICULTURE, FOOD AND ENVIRONMENT, LEXINGTON, KY, 40546

ID-36

Vegetable Production Guide for Commercial Growers, 2014-15

UK
UNIVERSITY OF
KENTUCKY
College of Agriculture,
Food and Environment



UKAg
EXTENSION

Agriculture and Natural Resources • Family and Consumer Sciences • 4-H Youth Development • Community and Economic Development

<http://www2.ca.uky.edu/agc/pubs/id/id36/id36.pdf>



UNIVERSITY OF ILLINOIS
EXTENSION



FERTILIZER: Tomatoes

The following fertilizer rates are to be used only as guidelines. Research at the University of Kentucky and at the University of Tennessee indicates that there is no yield increase from using more than 60 lb/A K_2O or 60 lb/A of P_2O_5 when soil test P and K levels are high.

Soil Test Results (lb/A)		Fertilizer Needed (lb/A)
Phosphorus		Phosphate (P_2O_5)
Low	<31	181-240
Medium	31-60	61-180
High	61-80	1-60
Very High	>80	0
Potassium		Potash (K_2O)
Low	<201	121-250
Medium	201-300	61-120
High	301-450	1-60
Very High	>450	0
Basal nitrogen where tomatoes:		N
1. follow grass-legume or legume sod		30
2. follow grass sod		50
3. are grown on continually cropped land		60

Supplemental applications: On bare ground plantings, apply an additional 30 lb of nitrogen/A as a sidedressing when the first fruits are golf-ball size. A second sidedress application of 30 lb N may also be desirable two or three weeks later, depending on the crop's growing condition. For plasticulture with drip on medium-textured soils, apply all recommended phosphorus and potassium requirements prior to laying plastic mulch. See fertigation table for N application rates.

FERTIGATION RECOMMENDATIONS (Nitrogen only): *Staked Tomatoes*

Based on a total season N recommendation of 150 lb actual N/A with 50 lb N/A applied preplant and the remaining N ($150 - 50 = 100$ lb) divided into equal amounts to be fertigated on a weekly basis ($100 \text{ lb} \div 10 \text{ weeks} = 10 \text{ lb of N per week}$). Both moderate (75 lb) and high (100 lb) fertigated N rates are provided below. For harvest seasons extending beyond 10 weeks from transplanting, a maintenance dose of 1 to 1.5 lb N (3 to 4.5 lb ammonium nitrate) per week is adequate. The doses listed for 1,000 plants are based on a plant population of 4,200 plants/A (i.e., rows on 6 ft centers in 5-row blocks and plants 18 in apart). IMPORTANT: If a UK soil test indicates your site is "at risk" for ripening disorders (Hartz ratio), you should alternate fertigations using ammonium or calcium nitrate with potassium nitrate fertigations (see Potassium Fertigation table).

Total Fertigated N Requirement¹	Actual N/wk (lb/A)	Ammonium Nitrate (lb/A/wk)	Ammonium Nitrate (lb/1,000 plants/wk)	Calcium Nitrate (lb/A/wk)	Calcium Nitrate (lb/1,000 plants/wk)
75 lb/A	7 lb 8 oz	22 lb 6 oz	5 lb 5 oz	48 lb 6 oz	11 lb 8 oz
100 lb/A	10 lb	30 lb	7 lb	64 lb 8 oz	15 lb 6 oz

¹ Fertigation can begin 10 to 14 days after transplanting and assumes 50 lb N/A was applied preplant and starter fertilizer was used.



FERTIGATION RECOMMENDATIONS (Nitrogen + Potassium): *Staked Tomatoes*

Potassium nitrate supplies both nitrogen and potassium and can be used as a substitute for ammonium or calcium nitrate. It is especially important to fertigate with a potassium source if a UK soil test indicates that your site is "at risk" for ripening disorders (Hartz ratio). Recommendations below are based on a total of 125 to 150 lb N/A for the season with 50 lb N/A applied preplant. The remaining $(125-50) = 75$ or $(150-50) = 100$ lb divided into equal amounts to be fertigated on a weekly basis for 10 weeks. This is either 7.5 or 10 lb of N/A/week. Both moderate (75 lb) and high (100 lb) fertigated N rates are given below. For harvest seasons extending beyond 10 weeks from transplanting, a maintenance dose of 1 to 1.5 lb N (11.5 lb potassium nitrate) per week is adequate. The doses listed for 1,000 plants are based on a plant population of 4,200 plants/A (i.e., rows 6 ft on center in 5-row blocks and plants 18 inches apart).

Total Fertigated N Requirement	Actual N/wk (lb/A)	Potassium Nitrate (lb/A/week)	Potassium Nitrate (lb/1,000 plants/wk)	Potassium Nitrate: K provided (lb/A/wk)	Potassium Nitrate: K provided (lb/1,000 plants/wk)
75 lb/A	7 lb 8 oz	57 lb 11 oz	13 lb 12 oz	25 lb 6 oz	6 lb 1 oz
100 lb/A	10 lb	76 lb 15 oz	18 lb 5 oz	33 lb 14 oz	8 lb 1 oz



Problems more commonly found in specialty crops

- Phosphorus scores too high
- Advising on foliar feeding
- Nutrients out of balance?
- “Disease” associated with fertility

Other differences

- Fertilizer materials used
- Timing of application
- Method of application
- Fertilizer input expenses are normally not that significant in the overall cost of production
- Excess fertilizer may result in problems beyond the simple waste of money

Tools in addition to soil testing utilized by Illinois specialty crop producers

- Tissue analysis
- Petiole sap meters
- Visual inspection

To reach us

Contacts	Contact information
Jeff Kindhart	jkindhar@illinois.edu
Rick Weinzierl	weinzier@illinois.edu