



UNIVERSITY OF ILLINOIS  
EXTENSION

**GROWING A NEW GENERATION  
OF ILLINOIS FRUIT AND VEGETABLE FARMERS**

# **PREVENTING VERTEBRATE PEST DAMAGE**

David Shiley, Extension Educator –  
Local Food Systems and Small Farms

**Illinois Migrant Council**



# Today's Objectives

- Understand wildlife nuisance management options for growers
- Understand legal implications of wildlife nuisance management
- Learn characteristics of common IL mammal species which can cause fruit and vegetable crop damage/and prevention of damage
- Understand integrated pest management for nuisance wildlife management

# What kind of animal is causing the damage?



## Is the damage seasonal or year round?

## Will the control program impact other species or the environment?



# Nuisance Wildlife Control Options

Population reduction

Habitat modification

Exclusion

Repellents

Frightening Devices

Integrate multiple strategies !!!!



# Population Reduction

Step one = permit

Department of Natural Resources in  
your state

Wildlife Protected by:

State Wildlife Codes

Federal Migratory Bird  
Treaty Act



Property owners do not own the wildlife !!!!

# Population Reduction

## Options:

- \*Seasonal Hunting

- \*Trapping

Live traps

Lethal

- \*Toxicants

Limited Use

# Population Reduction – Live Trapping

May be necessary for “problem animals”

## Problems with this strategy:

- \* Short-term fix
- \* Carrying Capacity

Amount of Food, Water, Shelter =  
number of animals area can support

- \* Release of live captured animals
- \* Very Expensive

# Population Reduction –

## Reduce total population

Reducing the population will reduce the effort required to control nuisance damage

Accomplished through recreational hunting and trapping.



# Principles of Habitat Modification

**Goal: Decrease the amount Food, Water, or Shelter**



**Examples:**

- \* Remove pruned branches/brush piles
- \* Mow tall grass or brushy areas

# Exclusion

## Fencing - protection

- \* Individual plants
- \* Planting areas



**Fencing = Expensive Option !!**

# Repellents

- \* Must be registered for use on species
- \* Regulated as a pesticide

## Types

- \* Taste repellents
- \* Olfactory - Area
- \* Tactile - birds



# Repellents: Positives and negatives

Pros:\* Can offer protection during periods of plant dormancy, or when damage needs to be limited during fruit set, ie sweet corn pollination.

Cons:\* Can be an expensive technique if year-round use is necessary

- \* Many products don't weather very well
- \* Most products can not be applied to fruit and vegetable crops



# Frightening Devices

- \*Visual

- \*Sound

Frightening techniques have limited success

Effectiveness increases with:

- \* Variation of strategies
- \* Persistence with tactics

## Biology:

Short life span = 12-15 months

Ave. = 4 to 6 months

3 - 6 litters per year



Home range = 10 acres

## Eastern Cottontail Rabbit



## Exclusion from trees:

- \* Wire mesh 1 inch or less
- \* Individually protect trees less than 6 years old with wire mesh tree guard at least 1.5 feet tall

# Repellents:

Products containing the fungicide Thiram, use limited to ornamentals



# Habitat Modification

Remove tall grass areas or keep away from tree plantings, orchards and vegetable crop fields.

Pick up and remove drops in the orchard, mow grass short between rows, remove vegetation in the row, eliminate connections to adjacent woodlands, pastures, and "old field" sites

# Population reduction

Trapping: Need permit to trap!!!!

Removal of problem animals in the winter can be effective in reducing urban populations.

Hunting:

Rabbits are an Upland Game species in Illinois, with an annual hunting season Nov. - Jan., an IL hunting license is required.

Removal of rabbits through trapping and hunting may reduce the current population, but may not reduce it the next season.



# White-tailed Deer

Average home range:

does 640 acres, bucks: 2000 – 3000 acres

Problems:

Browsing of trees and shrubs

Damage from antler rubbing

Damage to strawberries, melons and other crops





# Damage Identification



## Deer browse

Photo Credits: Dallas Virchow



## antler rubbing



UNIVERSITY OF ILLINOIS  
EXTENSION







Protect new trees in the landscape and orchard during the fall from antler rubbing

Use tree protectors such as Vexar, Tubex, plastic tree wrap or woven wire cylinders to protect new plantings.



# Exclusion from Area

## Fencing (Expensive)

Protecting a high value crop

Cost projected over life of fence

Low population design or temporary



Photo Credits: Dallas Virchow



# Deer fences need to be visible to animals

## Polytape materials

### Maintain 6 to 12 foot clear area between fence and outside edge area




Photo Credits: Dallas Virchow

\*Fence designs vary, dependent on population of deer and size of area to protect

Living with Wildlife in Illinois

<http://web.extension.illinois.edu/wildlife/>



\*Electrify sections of fencing as constructed



5.5 - 6 feet



Mike Roegge

OCT 25 2012







Mike Roegge

OCT 25 2012

# Deer Fence Netting

New material

Difficult for deer to see

Use techniques to increase visibility

Mylar tape, plastic flagging



# Taste Repellents

Usually high in cost with limitations

Examples (Active ingredients):

- \* Putrescent whole egg solids
- \* Thiram 
- \* Capsaicin best with an  
anti-transpirant






## Area (odor) repellents: active ingredients

- \* Tankage (Putrefied meat scraps)
- \* Edible animal protein
- \* Ammonium soaps of fatty acids
- \* Human Hair bags (3 feet apart)
- \* Bar deodorant soap (3 feet apart)

# Other Notes about Repellents

- Better suited for dormant trees and shrubs
- Repellents will most likely reduce, not eliminate damage 
- Effectiveness also dependent on size of deer population, and availability of other palatable foods

# Deer Population Management

## Recreational Hunting

Must reduce the number of females

More effective when adjacent  
landowners cooperate

## Nuisance population reduction

Deer Removal Permit required, based  
on damage, contact your DNR biologist  
Often only a temporary solution

# Habitat-vegetation management

Leave open areas between crop and woodland edges - 60 feet mowed

Use repellents and fencing near field edges

Forest management plans should include practices that promote abundance of young trees and shrubs for browse

Plant alternate food sources - food plots



# Voles

## Prairie and meadow voles

Tail is 2 times longer than the hind leg

Runways are on the soil surface

Damage is seen above the ground on tree trunk

## Pine voles

Tail is as long as the hind leg

Runways are usually underground

Damage is below ground as bark removed from roots and base of trunk

# Monitor the vole population - Apple Sign Test

30 monitoring stations per acre

3 rows of trees in orchard identified for test

10 stations per row , 30-40 feet apart

Shingles can be used but must be arched to allow voles to move under the shingle

5 days after placement, place  $\frac{1}{2}$  inch cube of apple under shingle

Check apple slice 24 hours later

Percent of apple slices with chewing evidence can give an estimate of potential damage

Monitoring stations help identify locations where vole damage is more likely



Conduct the test in the fall and spring and 21 to 30 days after rodenticide application

# Toxicants

Zinc phosphide- pelleted and grain bait formulation, more effective against meadow voles, but has high toxicity to other wildlife, therefore care must be taken to avoid killing non-target species



Anticoagulant baits - Chlorophacinone more effective against pine voles, requires multiple feedings



# Rodenticide application

Zinc phosphide can be broadcast applied into vegetated area, or used in a bait station within an orchard, but not on a continual basis as bait shyness may develop

Chlorophacinone can be used in a bait station over the winter to reduce the risk of re-infestation

Read and follow pesticide labels



# Non-chemical practices

Mow between the rows and trees within orchards on a regular basis, especially early fall

Herbicide strip or circle around fruit trees out to the drip line

Maintain a "clean" buffer around the orchard or crop field to limit movement of voles from adjacent habitat

Construct predator (hawk and owl) perches on the farm. 10 to 15 foot pole with "T" top

# Raccoons



Very common and adaptable mammal in Illinois

**Exclusion of raccoons from crop fields is the most effective control to reduce or prevent crop damage.**

Electric fencing can be used for exclusion

One wire 6 inches off the ground or a double wire with the second wire 6 inches above the first.

Fence chargers can be put on timers to turn on at dusk and off at sunrise

Fencing should be installed and charged prior to anticipated damage on sweet corn and melons





Inspect buildings and make repairs prior to problems. Do not leave feed sources outside and close doors and windows prior to sunset.

Electric fencing can be used to exclude raccoons and other mammals in areas where permanent structural changes can not be made.



# Damage to buildings from raccoons



ICWDM.ORG & USDA/WS



ICWDM.ORG & USDA/WS



ICWDM.ORG & USDA/WS

No toxicants, repellents or fumigants are registered for use to manage raccoons

Frightening:


Not effective for long periods of time to reduce damage



# Groundhog/Woodchuck

Small home range: 50 to 150 feet

## Control methods:

- Trapping 
- Fencing: Woven wire/electric combination
- Gas cartridge in burrow - CAUTION



Birds NOT  
protected by state  
and federal laws  
include .....

Starlings

Pigeons

House Sparrows



All other birds are protected by state  
and federal laws !!!!

# Exclusion from crops

- \*Exclude birds from area of damage with bird netting

- \*More successful if damage has just begun, feeding pattern not established

- \*Expensive and labor intensive, but may be necessary to reduce crop loss

Sapsucker damage can be extensive on select trees

Exclusion from feeding location on landscape trees.

May not be the best option for orchards, nurseries, etc. because damage is expanded



# Wildlife Damage Management Resources

\* Living With Wildlife in Illinois, U of I Extension

<http://web.extension.uiuc.edu/wildlife/>

\* Living With White-tailed Deer in Illinois, U of I Extension



<http://web.extension.illinois.edu/deer/>

\* The Internet Center for Wildlife Damage Management, University of Nebraska-Lincoln

<http://icwdm.org/>





# Reasons nuisance control programs fail

- Use of only one management strategy
- Not enough monitoring for damage
- Animals have become accustomed to using area
- Lack of "other" available food sources
- Lack of management plan or anticipation of seasonal damage
- Lack of persistence by landowner to change animal's behavior
- High population of problem animals



**Remember to use. . . .  
an Integrated Pest Management approach**

# To reach us

---

## Contacts

## Contact information

Rick Weinzierl

[weinzier@illinois.edu](mailto:weinzier@illinois.edu)

David Shiley

[dshiley@illinois.edu](mailto:dshiley@illinois.edu)

Other presenter

---